(12) STANDARD PATENT (19) AUSTRALIAN PATENT OFFICE

(11) Application No. AU 2002237586 B2

Document FP3 Appl. No. 10/690,639

(54)	Title Pharmaceutical composition of F(AB)2 fragments of antibodies and method for the prepartion thereof
(51)	International Palent Classification(s) C07K 16/06 (2006.01) C07K 16/18 (2006.01)
(21)	Application No: 2002237586 (22) Data of Filing: 2002.02.28
(87)	WIPO No: WO02/068475
(30)	Priority Data
(31)	Number (32) Date (33) Country 09/798,076 2001.03.05 US
(43) (43) (44)	Publication Date: 2002.09.12 Publication Journal Date: 2003.03.06 Accepted Journal Date: 2006.12.21
(71)	Applicant(s) Juan Lopez De Silanes;Rita Mancilla Nava;Jorgé Paniagua Solis
(72)	Inventor(s) Paniagua Solis, Jorge;Mancilla Nava, Rita;Lopez De Silanes, Juan
(74)	Agent / Attorney Cullen & Co, Level 26 239 George Street, Brisbane, QLD, 4000



LETTERS PATENT

STANDARD PATENT

2002237586

I, Fatima Beattie, the Commissioner of Patents, grant a Standard Patent with the following particulars:

Name and Address of Patentee(s):

Juan Lopez De Silanes

Lucema No. 7, Col., Juaiez Delegacion Cuaulitemoc, D.F. 06500, Mexico

Rita Mancilla Nava

Calzada de Tlalpan No. 4687, Colonia Toriello Guerra, Delegacion Tlalpan, D.F. 14950, Mexico.

Jorge Paniagua Solis

Amores No. 1304, Colonia del Valle, Delegación Benilo Juarez, D.F. 53100, Mexico

Name of Actual Inventor(s):

Paniagua Solis, Jorge; Mancilla Nava, Rita and Lopez De Silanes, Juan.

Title of Invention:

Pharmaceutical composition of F(AB)2 fragments of antibodies and method for the preparation

thereof

Term of Letters Patent:

Twenty years from 28 February 2002

Priority Details:

Number

Date

Filed wi

09/798,076

5 March 2001

US



Dated this 5th day of April 2007

PATENTS ACT 1990

Fatima Beattle Commissioner of Patents 15

The claims defining the invention are as follows:

- 1. A method for preparing a composition of F(ab')₂ antibody fragments that is substantially free of whole antibodies, the method comprising the steps of:
- (a) generating a source of antibodies by immunising animals with a complex antigenic mixture of animal venom, extracting blood plasma that is the source of antibodies and contacting with pepsin at a pH of 3.2 ± 0.2 under conditions to prepare an antibody digest containing $F(ab')_2$ fragments and being completely free of non-hydrolyzed antibodies;
- (b) treating said digested antibodies by two steps of ammonium sulphate precipitation, with at least one step at about 16% to about 22% weight by volume of ammonium sulphate for at least 30 minutes at a temperature of 55 °C \pm 4 °C , next cooling to approximately 8 °C \pm 4 °C for at least two hours;
 - (c) clarifying the solution by filtering at 12, 8, 4 or 0.22μ ;
- (d) subjecting the obtained solution to a second precipitation at about 32% to about 38% weight by volume of ammonium sulphate at about pH 6.8 \pm 0.5, next leaving the solution to settle for at least 12 hours in refrigeration;
- (e) centrifuging the resulting suspension to eliminate the supernatant and recover the paste of F(ah')₂ fragments precipitated in step (d);
- (f) subjecting the obtained paste of F(ab')₂ fragments to dialysis to eliminate salts and low weight components and
- 20 (g) passing the solution through a sterile filter
 - 2. The method of claim 1, wherein said venom is produced by a scorpion.
 - 3. The method of claim 1, wherein said venom is produced by a snake.
 - 4. The method of claim 1, wherein said venom is produced by a spider.
- 5. The method of claim 1 or claim 2, wherein the venom is selected from the group of scorpions of the taxonomic family Butidae and is selected one of the group consisting of:

 Centruroides elegans-elegans, Centruroides exilicanda, Centruroides infamatus-infamatus,

 Centruroides limpidus-tecomanus, Centruroides limpidus-limpidus, Centruroides suffusus.

- 6. The method of claim 1 or claim 3, wherein the venom is selected from one of the snake genera: Bothrops, Crotalus, Agkistrodon, Lachesis Sistrurus, or the species Micrurus nigrosinctus.
- The method of claim 1 or claim 4, wherein the venom is from the spider species: 7. Lactrodectus mactans.
 - The method of any one of claims 1 to 7, wherein the obtained composition of F(ab')2 8. fragments is free of albumin, free of whole antibodies and free of viruses and pyrogens.
 - A composition of F(ab')2 fragments having features a, b and c as follows: 9.
 - (a) the F(ab')₂ fragments are produced by the method of any one of claims 1 to 8,
 - (b) the F(ab')₂ fragments are free of whole antibodies, free of albumin, free of viral particles, and free of pyrogens,
- 15 (c) the F(ab')₂ fragments are polyclonal, and capable of neutralising all the antigenic determinants present in complex animal venoms utilized to generate the source of antibodies of step (a).
 - 10. The composition of claim 9 which includes a pharmaceutically acceptable carrier.

20

6 December 2006 Date: